arXiv:math/9908146v1 [math.CA] 27 Aug 1999

Inversion methods for finding differential equations for generalized Jacobi polynomials

J. Koekoek R. Koekoek

Abstract

We look for differential equations satisfied by the generalized Jacobi polynomials $\left\{P_n^{\alpha,\beta,M,N}(x)\right\}_{n=0}^{\infty}$ which are orthogonal on the interval $[-1,1]$ with respect to the weight function

$$\frac{\Gamma(\alpha+\beta+2)}{2^{\alpha+\beta+1}\Gamma(\alpha+1)\Gamma(\beta+1)}(1-x)^{\alpha}(1+x)^{\beta}+M\delta(x+1)+N\delta(x-1),$$

where $\alpha>-1$, $\beta>-1$, $M\geq 0$ and $N\geq 0$.

In order to find explicit formulas for the coefficients of these differential equations we have to solve systems of equations of the form

$$\sum_{i=1}^{\infty}A_i(x)D^iP_n^{(\alpha,\beta)}(x)=F_n(x),\ n=1,2,3,\ldots,$$

where the coefficients $\{A_i(x)\}_{i=1}^{\infty}$ are independent of n. This system of equations has a unique solution given by

$$A_i(x)=2^i\sum_{j=1}^{i}\frac{\alpha+\beta+2j+1}{(\alpha+\beta+j+1)_{i+1}}P_{i-j}^{(-\alpha-i-1,-\beta-i-1)}(x)F_j(x),\ i=1,2,3,\ldots.$$

This is a consequence of the Jacobi inversion formula

$$\sum_{k=j}^{i}\frac{\alpha+\beta+2k+1}{(\alpha+\beta+k+j+1)_{i-j+1}}\times$$
$$\times P_{i-k}^{(-\alpha-i-1,-\beta-i-1)}(x)P_{k-j}^{(\alpha+j,\beta+j)}(x)=\delta_{ij},\ j\leq i,\ i,j=0,1,2,\ldots,$$

which is proved in this report.

Keywords : Differential equations, Generalized Jacobi polynomials,
Inversion formulas.

1991 Mathematics Subject Classification : Primary 33C45 ; Secondary 34A35.

1 Introduction

In [13] T.H. Koornwinder introduced the generalized Jacobi polynomials $\left\{P_n^{\alpha,\beta,M,N}(x)\right\}_{n=0}^{\infty}$ which are orthogonal on the interval $[-1,1]$ with respect to the weight function

$$\frac{\Gamma(\alpha+\beta+2)}{2^{\alpha+\beta+1}\Gamma(\alpha+1)\Gamma(\beta+1)}(1-x)^{\alpha}(1+x)^{\beta}+M\delta(x+1)+N\delta(x-1),$$

where $\alpha>-1$, $\beta>-1$, $M\geq 0$ and $N\geq 0$. As a limit case he also found the generalized Laguerre polynomials $\left\{L_n^{\alpha,M}(x)\right\}_{n=0}^{\infty}$ which are orthogonal on the interval $[0,\infty)$ with respect to the weight function

$$\frac{1}{\Gamma(\alpha+1)}x^{\alpha}e^{-x}+M\delta(x),$$

where $\alpha>-1$ and $M\geq 0$. These generalized Jacobi polynomials and generalized Laguerre polynomials are related by the limit

$$L_n^{\alpha,M}(x)=\lim_{\beta\to\infty}P_n^{\alpha,\beta,0,M}\left(1-\frac{2x}{\beta}\right).$$

In [6] we proved that for $M>0$ the generalized Laguerre polynomials satisfy a unique differential equation of the form

$$M\sum_{i=0}^{\infty}a_i(x)y^{(i)}(x)+xy''(x)+(\alpha+1-x)y'(x)+ny(x)=0,$$

where $\{a_i(x)\}_{i=0}^{\infty}$ are continuous functions on the real line and $\{a_i(x)\}_{i=1}^{\infty}$ are independent of the degree n. In [1] H. Bavinck found a new method to obtain the main result of [6]. This inversion method was found in a similar way as was done in [4] in the case of generalizations of the Charlier polynomials. See also section 5 for more details. In [7] we used this inversion method to find all differential equations of the form

$$M\sum_{i=0}^{\infty}a_i(x)y^{(i)}(x)+N\sum_{i=0}^{\infty}b_i(x)y^{(i)}(x)+ \\ +MN\sum_{i=0}^{\infty}c_i(x)y^{(i)}(x)+xy''(x)+(\alpha+1-x)y'(x)+ny(x)=0, \qquad (1)$$

where the coefficients $\{a_i(x)\}_{i=1}^{\infty}$, $\{b_i(x)\}_{i=1}^{\infty}$ and $\{c_i(x)\}_{i=1}^{\infty}$ are independent of n and the coefficients $a_0(x)$, $b_0(x)$ and $c_0(x)$ are independent of x, satisfied by the Sobolev-type Laguerre polynomials $\left\{L_n^{\alpha,M,N}(x)\right\}_{n=0}^{\infty}$ which are orthogonal with respect to the inner product

$$<f,g>=\frac{1}{\Gamma(\alpha+1)}\int_0^{\infty}x^{\alpha}e^{-x}f(x)g(x)dx+Mf(0)g(0)+Nf'(0)g'(0),$$

where $\alpha>-1$, $M\geq 0$ and $N\geq 0$. These Sobolev-type Laguerre polynomials $\left\{L_n^{\alpha,M,N}(x)\right\}_{n=0}^{\infty}$ are generalizations of the generalized Laguerre polynomials $\left\{L_n^{\alpha,M}(x)\right\}_{n=0}^{\infty}$. In fact we have

$$L_n^{\alpha,M,0}(x)=L_n^{\alpha,M}(x) \text{ and } L_n^{\alpha,0}(x)=L_n^{(\alpha)}(x).$$

In this report we will prove an inversion formula involving the classical Jacobi polynomials which can be used to find differential equations of the form

$$M\sum_{i=0}^{\infty}a_i(x)y^{(i)}(x)+N\sum_{i=0}^{\infty}b_i(x)y^{(i)}(x)+MN\sum_{i=0}^{\infty}c_i(x)y^{(i)}(x)+ \\ +(1-x^2)y''(x)+\left[\beta-\alpha-(\alpha+\beta+2)x\right]y'(x)+n(n+\alpha+\beta+1)y(x)=0, \quad (2)$$

where the coefficients $\{a_i(x)\}_{i=1}^{\infty}$, $\{b_i(x)\}_{i=1}^{\infty}$ and $\{c_i(x)\}_{i=1}^{\infty}$ are independent of n and the coefficients $a_0(x)$, $b_0(x)$ and $c_0(x)$ are independent of x, satisfied by the generalized Jacobi polynomials $\left\{P_n^{\alpha,\beta,M,N}(x)\right\}_{n=0}^{\infty}$. As an example we apply the special case $\beta=\alpha$ of this inversion formula to solve the systems of equations obtained in [10].

The inversion formula for the Charlier polynomials obtained in [4] (see also section 5) was also used in [2] to find difference operators with Sobolev-type Charlier polynomials as eigenfunctions.

In [3] H. Bavinck and H. van Haeringen used similar inversion formulas to find difference equations for generalized Meixner polynomials.

2 The classical Laguerre and Jacobi polynomials

In this section we list the definitions and some properties of the classical Laguerre and Jacobi polynomials which we will use in this report. For details the reader is referred to [5], [12] and [16].

The classical Laguerre polynomials $\left\{L_n^{(\alpha)}(x)\right\}_{n=0}^{\infty}$ can be defined by

$$L_n^{(\alpha)}(x)=\sum_{k=0}^{n}(-1)^k\binom{n+\alpha}{n-k}\frac{x^k}{k!},\ n=0,1,2,\ldots \quad (3)$$

for all α. Their generating function is given by

$$(1-t)^{-\alpha-1}\exp\left(\frac{xt}{t-1}\right)=\sum_{n=0}^{\infty}L_n^{(\alpha)}(x)t^n \quad (4)$$

and for all $n\in\{0,1,2,\ldots\}$ we have

$$D^iL_n^{(\alpha)}(x)=(-1)^iL_{n-i}^{(\alpha+i)}(x),\ i=0,1,2,\ldots,n, \quad (5)$$

where $D=\dfrac{d}{dx}$ denotes the differentiation operator. The Laguerre polynomials satisfy the linear second order differential equation

$$xy''(x)+(\alpha+1-x)y'(x)+ny(x)=0. \quad (6)$$

It is well-known that

$$\frac{x^n}{n!}=\sum_{k=0}^{n}(-1)^k\binom{n+\alpha}{n-k}L_k^{(\alpha)}(x),\ n=0,1,2,\ldots. \quad (7)$$

This formula can easily be proved by using definition (3) and changing the order of summation as follows

$$\begin{aligned}\sum_{k=0}^{n}(-1)^k\binom{n+\alpha}{n-k}L_k^{(\alpha)}(x) &= \sum_{k=0}^{n}(-1)^k\binom{n+\alpha}{n-k}\sum_{j=0}^{k}(-1)^j\binom{k+\alpha}{k-j}\frac{x^j}{j!}\\ &= \sum_{j=0}^{n}\sum_{k=0}^{n-j}(-1)^k\binom{n+\alpha}{n-j-k}\binom{j+k+\alpha}{k}\frac{x^j}{j!}\\ &= \sum_{j=0}^{n}\binom{n+\alpha}{n-j}\frac{x^j}{j!}\sum_{k=0}^{n-j}(-1)^k\binom{n-j}{k}=\frac{x^n}{n!},\ n=0,1,2,\ldots.\end{aligned}$$

The classical Jacobi polynomials $\left\{P_n^{(\alpha,\beta)}(x)\right\}_{n=0}^{\infty}$ can be defined by

$$\begin{aligned}P_n^{(\alpha,\beta)}(x) &= \sum_{k=0}^{n}\frac{(n+\alpha+\beta+1)_k}{k!}\frac{(\alpha+k+1)_{n-k}}{(n-k)!}\left(\frac{x-1}{2}\right)^k,\ n=0,1,2,\ldots \quad (8)\\ &= (-1)^n\sum_{k=0}^{n}\frac{(-n-k-\alpha-\beta)_k}{k!}\frac{(-n-\alpha)_{n-k}}{(n-k)!}\left(\frac{x-1}{2}\right)^k,\ n=0,1,2,\ldots \quad (9)\\ &= 2^{-n}\sum_{k=0}^{n}\binom{n+\alpha}{n-k}\binom{n+\beta}{k}(x-1)^k(x+1)^{n-k},\ n=0,1,2,\ldots \quad (10)\end{aligned}$$

for all α and β. For all $n\in\{0,1,2,\ldots\}$ we have

$$D^iP_n^{(\alpha,\beta)}(x)=\frac{(n+\alpha+\beta+1)_i}{2^i}P_{n-i}^{(\alpha+i,\beta+i)}(x),\ i=0,1,2,\ldots,n. \quad (11)$$

The Jacobi polynomials satisfy the symmetry formula

$$P_n^{(\alpha,\beta)}(-x)=(-1)^nP_n^{(\beta,\alpha)}(x),\ n=0,1,2,\ldots \quad (12)$$

and the linear second order differential equation

$$(1-x^2)y''(x)+\left[\beta-\alpha-(\alpha+\beta+2)x\right]y'(x)+n(n+\alpha+\beta+1)y(x)=0. \quad (13)$$

Further we have for $\alpha+\beta+1>0$ (compare with [14], page 277, formula (30))

$$\left(\frac{1-x}{2}\right)^n=\sum_{k=0}^{n}\frac{(-n)_k(\alpha+k+1)_{n-k}(\alpha+\beta+2k+1)}{(\alpha+\beta+k+1)_{n+1}}P_k^{(\alpha,\beta)}(x),\ n=0,1,2,\ldots. \quad (14)$$

This formula is much less known than formula (7) for the Laguerre polynomials. However, the proof is quite similar. In section 6 we will prove a much more general formula.

We remark that (14) can be written in a more general form as

$$\begin{aligned}&\sum_{k=0}^{n}\frac{(-n)_k(\alpha+\beta+1)_k(\alpha+k+1)_{n-k}(\alpha+\beta+2k+1)}{\Gamma(\alpha+\beta+n+k+2)}P_k^{(\alpha,\beta)}(x)\\ &= \frac{1}{\Gamma(\alpha+\beta+1)}\left(\frac{1-x}{2}\right)^n,\ n=0,1,2,\ldots, \quad (15)\end{aligned}$$

which is valid for all α and β.

The Jacobi polynomials $\left\{P_n^{(\alpha,\beta)}(x)\right\}_{n=0}^{\infty}$ and the Laguerre polynomials $\left\{L_n^{(\alpha)}(x)\right\}_{n=0}^{\infty}$ are related by the limit

$$L_n^{(\alpha)}(x) = \lim_{\beta\to\infty} P_n^{(\alpha,\beta)}\left(1-\frac{2x}{\beta}\right). \tag{16}$$

We remark that if we replace x by $1-\dfrac{2x}{\beta}$ in (14), multiply by β^n and let β tend to infinity in the complex plane along the halfline where $\alpha+\beta$ is real and $\alpha+\beta+1>0$ we obtain (7) by using (16) and the fact that we have for all $n \in \{0,1,2,\ldots\}$

$$(-n)_k(\alpha+k+1)_{n-k} = (-1)^k\binom{n+\alpha}{n-k}n!, k=0,1,2,\ldots,n.$$

Finally we derive some formulas involving Jacobi polynomials which we will need in this report. By using definition (8) we easily obtain

$$\begin{aligned}
& nP_n^{(\alpha,\beta)}(x) - (n+\alpha)P_{n-1}^{(\alpha,\beta+1)}(x) \\
= {} & n\sum_{k=0}^{n}\frac{(n+\alpha+\beta+1)_k}{k!}\frac{(\alpha+k+1)_{n-k}}{(n-k)!}\left(\frac{x-1}{2}\right)^k + \\
& \qquad - (n+\alpha)\sum_{k=0}^{n-1}\frac{(n+\alpha+\beta+1)_k}{k!}\frac{(\alpha+k+1)_{n-k-1}}{(n-k-1)!}\left(\frac{x-1}{2}\right)^k \\
= {} & \sum_{k=0}^{n}\frac{(n+\alpha+\beta+1)_k}{k!}\frac{(\alpha+k+1)_{n-k}}{(n-k)!}\left[n-(n-k)\right]\left(\frac{x-1}{2}\right)^k \\
= {} & \sum_{k=0}^{n}\frac{(n+\alpha+\beta+1)_k}{k!}\frac{(\alpha+k+1)_{n-k}}{(n-k)!}(x-1)D\left(\frac{x-1}{2}\right)^k,\ n=1,2,3,\ldots.
\end{aligned}$$

Hence

$$(x-1)DP_n^{(\alpha,\beta)}(x) = nP_n^{(\alpha,\beta)}(x) - (n+\alpha)P_{n-1}^{(\alpha,\beta+1)}(x),\ n=1,2,3,\ldots. \tag{17}$$

By using the symmetry property (12) we also have

$$(x+1)DP_n^{(\alpha,\beta)}(x) = nP_n^{(\alpha,\beta)}(x) + (n+\beta)P_{n-1}^{(\alpha+1,\beta)}(x),\ n=1,2,3,\ldots. \tag{18}$$

Another easy consequence of definition (8) is

$$P_n^{(\alpha+1,\beta)}(x) - P_n^{(\alpha,\beta+1)}(x) = P_{n-1}^{(\alpha+1,\beta+1)}(x),\ n=1,2,3,\ldots. \tag{19}$$

If $\beta=\alpha$ we can combine the relations (17), (18) and (19) to find that

$$2xDP_n^{(\alpha,\alpha)}(x) = 2nP_n^{(\alpha,\alpha)}(x) + (n+\alpha)P_{n-2}^{(\alpha+1,\alpha+1)}(x),\ n=2,3,4,\ldots. \tag{20}$$

This formula turns out to be very useful in section 7. In that section we will also need the following well-known relation for ultraspherical polynomials (see for instance [16])

$$\begin{aligned}
&(n+2\alpha+1)(n+2\alpha+2)P_n^{(\alpha+1,\alpha+1)}(x) - (n+\alpha)(n+\alpha+1)P_{n-2}^{(\alpha+1,\alpha+1)}(x) \\
&\qquad = 2(n+\alpha+1)(2n+2\alpha+1)P_n^{(\alpha,\alpha)}(x),\ n=2,3,4,\ldots.
\end{aligned} \tag{21}$$

This formula can be proved by straightforward calculations as follows. As before we find by using definition (8)

$$\begin{aligned}
&(n+2\alpha+1)(n+2\alpha+2)P_n^{(\alpha+1,\alpha+1)}(x)-(n+\alpha)(n+\alpha+1)P_{n-2}^{(\alpha+1,\alpha+1)}(x)\\
=\ &(n+2\alpha+1)(n+2\alpha+2)\sum_{k=0}^{n}\frac{(n+2\alpha+3)_k}{k!}\frac{(\alpha+k+2)_{n-k}}{(n-k)!}\left(\frac{x-1}{2}\right)^k+\\
&\qquad -(n+\alpha)(n+\alpha+1)\sum_{k=0}^{n-2}\frac{(n+2\alpha+1)_k}{k!}\frac{(\alpha+k+2)_{n-k-2}}{(n-k-2)!}\left(\frac{x-1}{2}\right)^k\\
=\ &\sum_{k=0}^{n}\frac{(n+2\alpha+1)_k}{k!}\frac{(\alpha+k+2)_{n-k}}{(n-k)!}\times\\
&\qquad\times\left[(n+2\alpha+k+1)(n+2\alpha+k+2)-(n-k)(n-k-1)\right]\left(\frac{x-1}{2}\right)^k\\
=\ &2(2n+2\alpha+1)\sum_{k=0}^{n}\frac{(n+2\alpha+1)_k}{k!}\frac{(\alpha+k+1)_{n-k+1}}{(n-k)!}\left(\frac{x-1}{2}\right)^k\\
=\ &2(n+\alpha+1)(2n+2\alpha+1)P_n^{(\alpha,\alpha)}(x),\ n=2,3,4,\ldots.
\end{aligned}$$

Another useful relation involving ultraspherical polynomials is

$$(\alpha+1)P_n^{(\alpha+1,\alpha+1)}(x)-(n+\alpha+1)P_n^{(\alpha,\alpha)}(x)=\frac{1}{4}(n+\alpha+1)(1-x^2)P_{n-2}^{(\alpha+2,\alpha+2)}(x),\qquad(22)$$

which holds for $n=2,3,4,\ldots$. This can easily be shown by using definition (10) as follows

$$\begin{aligned}
&(\alpha+1)P_n^{(\alpha+1,\alpha+1)}(x)-(n+\alpha+1)P_n^{(\alpha,\alpha)}(x)\\
=\ &2^{-n}\sum_{k=0}^{n}\left[(\alpha+1)\binom{n+\alpha+1}{n-k}\binom{n+\alpha+1}{k}+\right.\\
&\qquad\left.-(n+\alpha+1)\binom{n+\alpha}{n-k}\binom{n+\alpha}{k}\right](x-1)^k(x+1)^{n-k}\\
=\ &-(n+\alpha+1)2^{-n}\sum_{k=1}^{n-1}\binom{n+\alpha}{n-k-1}\binom{n+\alpha}{k-1}(x-1)^k(x+1)^{n-k}\\
=\ &(n+\alpha+1)(1-x^2)2^{-n}\sum_{k=0}^{n-2}\binom{n+\alpha}{n-k-2}\binom{n+\alpha}{k}(x-1)^k(x+1)^{n-k-2}\\
=\ &\frac{1}{4}(n+\alpha+1)(1-x^2)P_{n-2}^{(\alpha+2,\alpha+2)}(x),\ n=2,3,4,\ldots.
\end{aligned}$$

3 Some summation formulas

In this section we will derive some summation formulas which will be used in this report. We start with a partial sum of a ${}_1F_0$ hypergeometric series at the point 1. In fact we have

$$\sum_{k=0}^{n}\binom{a+k}{k}=\sum_{k=0}^{n}\frac{(a+1)_k}{k!}=\frac{(a+2)_n}{n!}=\binom{n+a+1}{n},\ n=0,1,2,\ldots.\qquad(23)$$

This formula appeared as a special case of lemma 4 in [7]. Further we have the well-known Vandermonde summation formula

$$ {}_2F_1\left(\left.\begin{array}{c} -n, b \\ c \end{array}\right| 1\right) = \frac{(c-b)_n}{(c)_n},\ (c)_n \neq 0,\ n = 0, 1, 2, \ldots. \tag{24} $$

By using (24) we find that

$$ \begin{aligned} & \sum_{k=0}^{n} \frac{(-n)_k (b)_k}{(b+n+1)_k k!} (b+2k) \\ = \;& b\, {}_2F_1\left(\left.\begin{array}{c} -n, b+1 \\ b+n+1 \end{array}\right| 1\right) - \frac{nb}{b+n+1}\, {}_2F_1\left(\left.\begin{array}{c} -n+1, b+1 \\ b+n+2 \end{array}\right| 1\right) \\ = \;& b\, \frac{(n)_n}{(b+n+1)_n} - \frac{nb}{b+n+1} \frac{(n+1)_{n-1}}{(b+n+2)_{n-1}} = 0,\ (b+n+1)_n \neq 0,\ n = 1, 2, 3, \ldots. \end{aligned} \tag{25} $$

We remark that this can be written in a more general form as

$$ \sum_{k=0}^{n} \frac{(-n)_k (b)_k}{\Gamma(b+n+k+1) k!} (b+2k) = 0,\ n = 1, 2, 3, \ldots, \tag{26} $$

which is valid for all b. This formula will be used in section 6.

We also have the well-known Saalschütz summation formula

$$ {}_3F_2\left(\left.\begin{array}{c} -n, a, n+b+c-1 \\ b, a+c \end{array}\right| 1\right) = \frac{(b-a)_n (c)_n}{(b)_n (a+c)_n},\ (b)_n (a+c)_n \neq 0,\ n = 0, 1, 2, \ldots. \tag{27} $$

Note that this implies that

$$ \frac{(c)_n}{(b)_n} = {}_3F_2\left(\left.\begin{array}{c} -n, \frac{1}{2}(b-c), n+b+c-1 \\ b, \frac{1}{2}(b+c) \end{array}\right| 1\right),\ (b)_n ((b+c)/2)_n \neq 0,\ n = 0, 1, 2, \ldots. \tag{28} $$

This can be used to show that for $(\frac{1}{2}b + \frac{1}{2})_n (b+n+1)_n (b-c+1)_n \neq 0$

$$ \sum_{k=0}^{n} \frac{(-n)_k (b)_k (c)_k}{(b+n+1)_k (b-c+1)_k k!} (b+2k) = \frac{(b)_{n+1} (\frac{1}{2}b - c + \frac{1}{2})_n}{(b-c+1)_n (\frac{1}{2}b + \frac{1}{2})_n},\ n = 0, 1, 2, \ldots. \tag{29} $$

So we suppose that $n \in \{0, 1, 2, \ldots\}$ and that $(\frac{1}{2}b + \frac{1}{2})_n (b+n+1)_n (b-c+1)_n \neq 0$. Then we use (28) and change the order of summation to obtain

$$ \begin{aligned} & \sum_{k=0}^{n} \frac{(-n)_k (b)_k (c)_k}{(b+n+1)_k (b-c+1)_k k!} (b+2k) \\ = \;& \sum_{k=0}^{n} \frac{(-n)_k (b)_k}{(b+n+1)_k k!} (b+2k)\, {}_3F_2\left(\left.\begin{array}{c} -k, \frac{1}{2}b - c + \frac{1}{2}, b+k \\ b-c+1, \frac{1}{2}b + \frac{1}{2} \end{array}\right| 1\right) \\ = \;& \sum_{k=0}^{n} \frac{(-n)_k (b)_k}{(b+n+1)_k k!} (b+2k) \sum_{m=0}^{k} \frac{(-k)_m (\frac{1}{2}b - c + \frac{1}{2})_m (b+k)_m}{(b-c+1)_m (\frac{1}{2}b + \frac{1}{2})_m m!} \\ = \;& \sum_{m=0}^{n} \sum_{k=0}^{n-m} \frac{(-n)_{m+k} (-m-k)_m (b)_{2m+k} (\frac{1}{2}b - c + \frac{1}{2})_m}{(b+n+1)_{m+k} (b-c+1)_m (\frac{1}{2}b + \frac{1}{2})_m (m+k)!\, m!} (b+2m+2k) \\ = \;& \sum_{m=0}^{n} (-1)^m \frac{(-n)_m (b)_{2m} (\frac{1}{2}b - c + \frac{1}{2})_m}{(b+n+1)_m (b-c+1)_m (\frac{1}{2}b + \frac{1}{2})_m m!} \times \\ & \qquad \times \sum_{k=0}^{n-m} \frac{(-n+m)_k (b+2m)_k}{(b+n+m+1)_k k!} (b+2m+2k). \end{aligned} $$

Hence, by using (25) we conclude that

$$\begin{aligned}
&\sum_{k=0}^{n}\frac{(-n)_k(b)_k(c)_k}{(b+n+1)_k(b-c+1)_k k!}(b+2k)\\
=\;& (-1)^n\frac{(-n)_n(b)_{2n}(\frac{1}{2}b-c+\frac{1}{2})_n}{(b+n+1)_n(b-c+1)_n(\frac{1}{2}b+\frac{1}{2})_n n!}(b+2n)\\
=\;& \frac{(b)_{2n+1}(\frac{1}{2}b-c+\frac{1}{2})_n}{(b+n+1)_n(b-c+1)_n(\frac{1}{2}b+\frac{1}{2})_n}=\frac{(b)_{n+1}(\frac{1}{2}b-c+\frac{1}{2})_n}{(b-c+1)_n(\frac{1}{2}b+\frac{1}{2})_n},\ n=0,1,2,\ldots,
\end{aligned}$$

which proves (29).

4 The systems of equations

Let $\alpha>-1$. The Sobolev-type Laguerre polynomials $\left\{L_n^{\alpha,M,N}(x)\right\}_{n=0}^{\infty}$ can be written as

$$L_n^{\alpha,M,N}(x)=A_0L_n^{(\alpha)}(x)+A_1DL_n^{(\alpha)}(x)+A_2D^2L_n^{(\alpha)}(x),\ n=0,1,2,\ldots,$$

where the coefficients A_0, A_1 and A_2 are given by

$$\left\{\begin{aligned}
A_0&=1+M\binom{n+\alpha}{n-1}+\frac{n(\alpha+2)-(\alpha+1)}{(\alpha+1)(\alpha+3)}N\binom{n+\alpha}{n-2}+\\
&\qquad\qquad+\frac{MN}{(\alpha+1)(\alpha+2)}\binom{n+\alpha}{n-1}\binom{n+\alpha+1}{n-2}\\
A_1&=M\binom{n+\alpha}{n}+\frac{n-1}{\alpha+1}N\binom{n+\alpha}{n-1}+\frac{2MN}{(\alpha+1)^2}\binom{n+\alpha}{n}\binom{n+\alpha+1}{n-2}\\
A_2&=\frac{N}{\alpha+1}\binom{n+\alpha}{n-1}+\frac{MN}{(\alpha+1)^2}\binom{n+\alpha}{n}\binom{n+\alpha+1}{n-1}.
\end{aligned}\right.$$

For details concerning these Sobolev-type Laguerre polynomials and their definition the reader is referred to [8] and [11]. Since the classical Laguerre polynomials $\left\{L_n^{(\alpha)}(x)\right\}_{n=0}^{\infty}$ satisfy the differential equation (6) it is quite reasonable to look for differential equations of the form (1) for these Sobolev-type Laguerre polynomials $\left\{L_n^{\alpha,M,N}(x)\right\}_{n=0}^{\infty}$ in view of this definition and the fact that $L_n^{\alpha,0,0}(x)=L_n^{(\alpha)}(x)$. In [7] it is shown that this leads to eight systems of equations for the coefficients $\{a_i(x)\}_{i=0}^{\infty}$, $\{b_i(x)\}_{i=0}^{\infty}$ and $\{c_i(x)\}_{i=0}^{\infty}$. In order to find these coefficients we have to solve systems of equations which are of the form

$$\sum_{i=1}^{\infty}A_i(x)D^{i+k}L_n^{(\alpha)}(x)=F_n(x),\ n=k+1,k+2,k+3,\ldots,$$

where $k\in\{0,1,2,\ldots\}$ and the coefficients $\{A_i(x)\}_{i=1}^{\infty}$ are independent of n. In [7] it is pointed out that this system of equations has a unique solution given by

$$A_i(x)=(-1)^{i+k}\sum_{j=1}^{i}L_{i-j}^{(-\alpha-i-k-1)}(-x)F_{j+k}(x),\ i=1,2,3,\ldots.$$

This is an easy consequence of the Laguerre inversion formula

$$\sum_{k=j}^{i} L_{i-k}^{(-\alpha-i-1)}(-x) L_{k-j}^{(\alpha+j)}(x) = \delta_{ij},\ j \leq i,\ i,j = 0,1,2,\ldots, \tag{30}$$

which was found by H. Bavinck in [1]. For more details the reader is referred to [1] and [7]. See also section 5 of this report.

Now we take $\alpha > -1$ and $\beta > -1$. The generalized Jacobi polynomials $\left\{P_n^{\alpha,\beta,M,N}(x)\right\}_{n=0}^{\infty}$ can be written as

$$P_n^{\alpha,\beta,M,N}(x) = A_0 P_n^{(\alpha,\beta)}(x) + \left[A_1(1-x) - A_2(1+x)\right] DP_n^{(\alpha,\beta)}(x),\ n = 0,1,2,\ldots,$$

where the coefficients A_0, A_1 and A_2 are given by

$$\begin{cases} A_0 = 1 + M\dfrac{\dbinom{n+\beta}{n-1}\dbinom{n+\alpha+\beta+1}{n}}{\dbinom{n+\alpha}{n}} + N\dfrac{\dbinom{n+\alpha}{n-1}\dbinom{n+\alpha+\beta+1}{n}}{\dbinom{n+\beta}{n}} + \\ \qquad\qquad + MN\dfrac{(\alpha+\beta+2)^2}{(\alpha+1)(\beta+1)}\dbinom{n+\alpha+\beta+1}{n-1}^2 \\ A_1 = \dfrac{M}{\alpha+\beta+1}\dfrac{\dbinom{n+\beta}{n}\dbinom{n+\alpha+\beta}{n}}{\dbinom{n+\alpha}{n}} + \dfrac{MN}{\alpha+1}\dbinom{n+\alpha+\beta}{n-1}\dbinom{n+\alpha+\beta+1}{n} \\ A_2 = \dfrac{N}{\alpha+\beta+1}\dfrac{\dbinom{n+\alpha}{n}\dbinom{n+\alpha+\beta}{n}}{\dbinom{n+\beta}{n}} + \dfrac{MN}{\beta+1}\dbinom{n+\alpha+\beta}{n-1}\dbinom{n+\alpha+\beta+1}{n}. \end{cases}$$

Here we used the same definition as in [13], but in a slightly different notation. The case $\alpha + \beta + 1 = 0$ must be understood by continuity. In view of this definition and the fact that the classical Jacobi polynomials $\left\{P_n^{(\alpha,\beta)}(x)\right\}_{n=0}^{\infty}$ satisfy the differential equation (13) it is quite natural to look for differential equations of the form (2) satisfied by these generalized Jacobi polynomials as was already pointed out in [9]. Again this leads to eight systems of equations for the coefficients $\{a_i(x)\}_{i=0}^{\infty}$, $\{b_i(x)\}_{i=0}^{\infty}$ and $\{c_i(x)\}_{i=0}^{\infty}$. In order to find these coefficients we have to solve systems of equations which are of the form

$$\sum_{i=1}^{\infty} A_i(x) D^i P_n^{(\alpha,\beta)}(x) = F_n(x),\ n = 1,2,3,\ldots,$$

where the coefficients $\{A_i(x)\}_{i=1}^{\infty}$ are independent of n. This system of equations has a unique solution given by

$$A_i(x) = 2^i \sum_{j=1}^{i} \frac{\alpha+\beta+2j+1}{(\alpha+\beta+j+1)_{i+1}} P_{i-j}^{(-\alpha-i-1,-\beta-i-1)}(x) F_j(x),\ i = 1,2,3,\ldots.$$

This is a consequence of the Jacobi inversion formula

$$\sum_{k=j}^{i} \frac{\alpha+\beta+2k+1}{(\alpha+\beta+k+j+1)_{i-j+1}} \times \\ \times P_{i-k}^{(-\alpha-i-1,-\beta-i-1)}(x) P_{k-j}^{(\alpha+j,\beta+j)}(x) = \delta_{ij},\ j \leq i,\ i,j = 0,1,2,\ldots, \tag{31}$$

which will be proved in this report. Again, the case $\alpha+\beta+1=0$ must be understood by continuity. We remark that if we replace x by $1-\dfrac{2x}{\beta}$ in (31), multiply by β^{i-j} and let β tend to infinity along the positive real axis we obtain the Laguerre inversion formula (30) by using (16).

In [10] we found all differential equations of the form

$$M\sum_{i=0}^{\infty} a_i(x)y^{(i)}(x) + (1-x^2)y''(x) - 2(\alpha+1)xy'(x) + n(n+2\alpha+1)y(x) = 0, \tag{32}$$

where $\{a_i(x)\}_{i=0}^{\infty}$ are continuous functions on the real line and $\{a_i(x)\}_{i=1}^{\infty}$ are independent of n, satisfied by the symmetric generalized ultraspherical polynomials $\left\{P_n^{\alpha,\alpha,M,M}(x)\right\}_{n=0}^{\infty}$ defined by

$$P_n^{\alpha,\alpha,M,M}(x) = C_0 P_n^{(\alpha,\alpha)}(x) - C_1 x D P_n^{(\alpha,\alpha)}(x),\ n = 0,1,2,\ldots,$$

where

$$\begin{cases} C_0 = 1 + \dfrac{2Mn}{\alpha+1}\dbinom{n+2\alpha+1}{n} + 4M^2\dbinom{n+2\alpha+1}{n-1}^2 \\[2ex] C_1 = \dfrac{2M}{2\alpha+1}\dbinom{n+2\alpha}{n} + \dfrac{2M^2}{\alpha+1}\dbinom{n+2\alpha}{n-1}\dbinom{n+2\alpha+1}{n}. \end{cases}$$

We remark that these polynomials form a special case ($\beta=\alpha$ and $N=M$) of the generalized Jacobi polynomials $\left\{P_n^{\alpha,\beta,M,N}(x)\right\}_{n=0}^{\infty}$, but the differential equation (32) has a very special form without a M^2-part. This differential equation will appear not to be a special case of the differential equation of the form (2) for the generalized Jacobi polynomials, since the MN-part will not vanish if we take $\beta=\alpha$ and $N=M$. We aim to give a proof of this in a future publication. In section 7 of this report we will apply the special case $\beta=\alpha$ of the Jacobi inversion formula (31) to solve the systems of equations obtained in [10] as an example.

5 The inversion formulas

In [4] H. Bavinck and R. Koekoek found the following inversion formula involving Charlier polynomials

$$\sum_{k=j}^{i} C_{i-k}^{(-a)}(-x) C_{k-j}^{(a)}(x) = \delta_{ij},\ j \leq i,\ i,j = 0,1,2,\ldots. \tag{33}$$

This formula is an easy consequence of the generating function (see for instance [12])

$$e^{-at}(1+t)^x = \sum_{n=0}^{\infty} C_n^{(a)}(x)t^n.$$

In fact we have

$$\begin{aligned} 1 &= e^{-at}(1+t)^x e^{at}(1+t)^{-x} = \sum_{k=0}^{\infty} C_k^{(a)}(x)t^k \sum_{m=0}^{\infty} C_m^{(-a)}(-x)t^m \\ &= \sum_{n=0}^{\infty}\left(\sum_{k=0}^{n} C_k^{(a)}(x)C_{n-k}^{(-a)}(-x)\right)t^n. \end{aligned}$$

Hence

$$\sum_{k=0}^{n} C_k^{(a)}(x)C_{n-k}^{(-a)}(-x) = \begin{cases} 1, & n=0 \\ \\ 0, & n=1,2,3,\ldots. \end{cases}$$

Now (33) easily follows by taking $n=i-j$ and shifting the summation index. This formula was also used in [2] to find difference operators with Sobolev-type Charlier polynomials as eigenfunctions. In [3] a similar formula involving Meixner polynomials was used to find difference equations for generalized Meixner polynomials.

Formula (33) can be interpreted as follows. If we define the matrix $T=(t_{ij})_{i,j=0}^n$ with entries

$$t_{ij} = \begin{cases} C_{i-j}^{(a)}(x), & j \le i \\ \\ 0, & j > i, \end{cases}$$

then this matrix T is a triangular matrix with determinant 1 and the inverse U of this matrix is given by $T^{-1}=U=(u_{ij})_{i,j=0}^n$ with entries

$$u_{ij} = \begin{cases} C_{i-j}^{(-a)}(-x), & j \le i \\ \\ 0, & j > i. \end{cases}$$

Therefore we call (33) an inversion formula.

In the same way we find by using the generating function (4) for the Laguerre polynomials

$$\sum_{k=j}^{i} L_{i-k}^{(\alpha)}(x)L_{k-j}^{(-\alpha-2)}(-x) = \delta_{ij},\ j \le i,\ i,j=0,1,2,\ldots.$$

However, this formula cannot be used to solve systems of equations of the form

$$\sum_{i=1}^{\infty} A_i(x)D^iL_n^{(\alpha)}(x) = F_n(x),\ n=1,2,3,\ldots$$

in view of the parametershift in (5).

In [1] H. Bavinck used a slightly different method to find the Laguerre inversion formula (30) from the generating function (4) for the Laguerre polynomials. In fact we have

$$\begin{aligned} (1-t)^{i-j-1} &= (1-t)^{-\alpha-j-1}\exp\left(\frac{xt}{t-1}\right)(1-t)^{\alpha+i}\exp\left(\frac{-xt}{t-1}\right) \\ &= \sum_{k=0}^{\infty} L_k^{(\alpha+j)}(x)t^k \sum_{m=0}^{\infty} L_m^{(-\alpha-i-1)}(-x)t^m \\ &= \sum_{n=0}^{\infty}\left(\sum_{k=0}^{n} L_k^{(\alpha+j)}(x)L_{n-k}^{(-\alpha-i-1)}(-x)\right)t^n. \end{aligned}$$

This implies, by comparing the coefficients of t^{i-j} on both sides, that

$$\sum_{k=0}^{i-j} L_k^{(\alpha+j)}(x) L_{i-j-k}^{(-\alpha-i-1)}(-x) = \delta_{ij},\ j \le i,\ i,j = 0,1,2,\ldots,$$

which is equivalent to (30).

Formula (30) can be interpreted as follows. If we define the matrix $T = (t_{ij})_{i,j=0}^n$ with entries

$$t_{ij} = \begin{cases} L_{i-j}^{(\alpha+j)}(x), & j \le i \\ \\ 0, & j > i, \end{cases}$$

then this matrix T is a triangular matrix with determinant 1 and the inverse U of this matrix is given by $T^{-1} = U = (u_{ij})_{i,j=0}^n$ with entries

$$u_{ij} = \begin{cases} L_{i-j}^{(-\alpha-i-1)}(-x), & j \le i \\ \\ 0, & j > i. \end{cases}$$

It is possible to generalize the Laguerre inversion formula to

$$\sum_{k=0}^{n} L_k^{(\alpha+p_n)}(x) L_{n-k}^{(-\alpha-q_n)}(-x) = \frac{(p_n - q_n + 2)_n}{n!},\ n = 0,1,2,\ldots, \tag{34}$$

where p_n and q_n are arbitrary and even may depend on n. In order to have an inversion formula we have to choose p_n and q_n such that

$$(p_n - q_n + 2)_n = 0,\ n = 1,2,3,\ldots,$$

hence

$$p_n - q_n \in \{-n-1, -n, \ldots, -3, -2\},\ n = 1,2,3,\ldots.$$

Note that the endpoint-cases $p_n - q_n = -2$ and $p_n - q_n = -n-1$ correspond to the earlier mentioned inversion formulas.

To prove (34) we use (4) to obtain

$$L_n^{(\alpha)}(x) = \frac{1}{n!}\, D_t^n \left[(1-t)^{-\alpha-1} \exp\left(\frac{xt}{t-1}\right)\right]\Bigg|_{t=0},\ n = 0,1,2,\ldots,$$

where $D_t = \dfrac{d}{dt}$ denotes differentiation with respect to t. Hence by using Leibniz' rule we find

$$\begin{aligned} &\sum_{k=0}^{n} L_k^{(\alpha+p_n)}(x) L_{n-k}^{(-\alpha-q_n)}(-x) \\ = &\sum_{k=0}^{n} \frac{1}{k!}\, D_t^k \left[(1-t)^{-\alpha-p_n-1} \exp\left(\frac{xt}{t-1}\right)\right]\Bigg|_{t=0} \times \\ &\qquad \times \frac{1}{(n-k)!}\, D_t^{n-k} \left[(1-t)^{\alpha+q_n-1} \exp\left(\frac{-xt}{t-1}\right)\right]\Bigg|_{t=0} \\ = &\frac{1}{n!}\, D_t^n \left[(1-t)^{q_n-p_n-2}\right]\Big|_{t=0} = \frac{(p_n - q_n + 2)_n}{n!},\ n = 0,1,2,\ldots, \end{aligned}$$

which proves (34).

In case of the Jacobi polynomials the above methods seem not to be applicable. However, in the next section we will give a proof of the Jacobi inversion formula (31).

6 Proof of the Jacobi inversion formula

In this section we will prove that

$$\begin{aligned}&\sum_{k=0}^{n}\frac{(\alpha+\beta+2k+1)(\alpha+\beta+1)_k}{\Gamma(\alpha+\beta+n+k+2)}P_k^{(\alpha,\beta)}(x)P_{n-k}^{(-n-\alpha-1,-n-\beta-1)}(y)\\ = \;&\frac{1}{\Gamma(\alpha+\beta+1)}\frac{1}{n!}\left(\frac{x-y}{2}\right)^n,\ n=0,1,2,\ldots,\end{aligned}\tag{35}$$

which holds for all α and β.

Note that (15) is a special case of (35) since

$$\begin{aligned}P_{n-k}^{(-n-\alpha-1,-n-\beta-1)}(1) &= \frac{(-n-\alpha)_{n-k}}{(n-k)!}=(-1)^{n-k}\frac{(\alpha+k+1)_{n-k}}{(n-k)!}\\ &= \frac{(-1)^n}{n!}(-n)_k(\alpha+k+1)_{n-k},\ k=0,1,2,\ldots,n\end{aligned}$$

for all $n\in\{0,1,2,\ldots\}$.

By taking $y=x$ in (35) we easily obtain

$$\begin{aligned}&\sum_{k=0}^{n}\frac{(\alpha+\beta+2k+1)(\alpha+\beta+1)_k}{\Gamma(\alpha+\beta+n+k+2)}\times\\ &\qquad\times P_k^{(\alpha,\beta)}(x)P_{n-k}^{(-n-\alpha-1,-n-\beta-1)}(x)=\begin{cases}\dfrac{1}{\Gamma(\alpha+\beta+1)}, & n=0\\ 0, & n=1,2,3,\ldots\end{cases}\end{aligned}\tag{36}$$

for all α and β. If we take $n=i-j$ in (36) and shift the summation index we find

$$\begin{aligned}&\sum_{k=j}^{i}\frac{(\alpha+\beta+2k-2j+1)(\alpha+\beta+1)_{k-j}}{\Gamma(\alpha+\beta+i-2j+k+2)}\times\\ &\qquad\times P_{i-k}^{(-i+j-\alpha-1,-i+j-\beta-1)}(x)P_{k-j}^{(\alpha,\beta)}(x)=\frac{\delta_{ij}}{\Gamma(\alpha+\beta+1)},\ j\le i,\ i,j=0,1,2,\ldots.\end{aligned}$$

For α and β real with $\alpha+\beta+1>-1$ we now obtain (31) by shifting both α and β by j.

Note that (35) for $y=-x$ in a similar way leads to

$$\begin{aligned}&\sum_{k=j}^{i}\frac{\alpha+\beta+2k+1}{(\alpha+\beta+k+j+1)_{i-j+1}}\times\\ &\qquad\times P_{i-k}^{(-\alpha-i-1,-\beta-i-1)}(-x)P_{k-j}^{(\alpha+j,\beta+j)}(x)=\frac{x^{i-j}}{(i-j)!},\ j\le i,\ i,j=0,1,2,\ldots,\end{aligned}\tag{37}$$

which will also be used in the next section.

In order to prove (35) we start with the left-hand side, apply definition (8) to $P_k^{(\alpha,\beta)}(x)$ and definition (9) to $P_{n-k}^{(-n-\alpha-1,-n-\beta-1)}(y)$ and change the order of summation to obtain

$$\sum_{k=0}^{n}\frac{(\alpha+\beta+2k+1)(\alpha+\beta+1)_k}{\Gamma(\alpha+\beta+n+k+2)}P_k^{(\alpha,\beta)}(x)P_{n-k}^{(-n-\alpha-1,-n-\beta-1)}(y)$$

$$\begin{aligned}
&= \sum_{k=0}^{n}\sum_{i=0}^{k}\sum_{j=0}^{n-k}(-1)^{n-k}\frac{(\alpha+\beta+2k+1)(\alpha+\beta+1)_k}{\Gamma(\alpha+\beta+n+k+2)}\frac{(\alpha+\beta+k+1)_i}{i!}\frac{(\alpha+i+1)_{k-i}}{(k-i)!}\\
&\qquad\times\frac{(\alpha+\beta+n+k-j+2)_j}{j!}\frac{(\alpha+k+1)_{n-k-j}}{(n-k-j)!}\left(\frac{x-1}{2}\right)^i\left(\frac{y-1}{2}\right)^j\\
&= \sum_{i=0}^{n}\sum_{k=i}^{n}\sum_{j=0}^{n-k}(-1)^{n-k}\times\\
&\qquad\times\frac{(\alpha+\beta+2k+1)(\alpha+\beta+1)_{i+k}(\alpha+i+1)_{n-i-j}}{\Gamma(\alpha+\beta+n+k-j+2)\,i!\,(k-i)!\,j!\,(n-k-j)!}\left(\frac{x-1}{2}\right)^i\left(\frac{y-1}{2}\right)^j\\
&= \sum_{i=0}^{n}\sum_{k=0}^{n-i}\sum_{j=0}^{n-i-k}(-1)^{n-i-k}\times\\
&\qquad\times\frac{(\alpha+\beta+2i+2k+1)(\alpha+\beta+1)_{2i+k}(\alpha+i+1)_{n-i-j}}{\Gamma(\alpha+\beta+n+i+k-j+2)\,i!\,k!\,j!\,(n-i-k-j)!}\left(\frac{x-1}{2}\right)^i\left(\frac{y-1}{2}\right)^j\\
&= \sum_{i=0}^{n}\sum_{j=0}^{n-i}\sum_{k=0}^{n-i-j}(-1)^{n-i}\left(\frac{x-1}{2}\right)^i\left(\frac{y-1}{2}\right)^j\times\\
&\qquad\times\frac{(\alpha+\beta+2i+2k+1)(\alpha+\beta+1)_{2i+k}(\alpha+i+1)_{n-i-j}(-n+i+j)_k}{\Gamma(\alpha+\beta+n+i-j+k+2)\,i!\,j!\,k!\,(n-i-j)!}\\
&= \sum_{i=0}^{n}\sum_{j=0}^{n-i}(-1)^{n-i}\frac{(\alpha+\beta+1)_{2i}(\alpha+i+1)_{n-i-j}}{i!\,j!\,(n-i-j)!}\left(\frac{x-1}{2}\right)^i\left(\frac{y-1}{2}\right)^j\times\\
&\qquad\times\sum_{k=0}^{n-i-j}\frac{(-n+i+j)_k(\alpha+\beta+2i+1)_k}{\Gamma(\alpha+\beta+n+i-j+k+2)k!}(\alpha+\beta+2i+2k+1),\ n=0,1,2,\ldots.
\end{aligned}$$

Now we use (26) to obtain

$$\begin{aligned}
&\sum_{k=0}^{n}\frac{(\alpha+\beta+2k+1)(\alpha+\beta+1)_k}{\Gamma(\alpha+\beta+n+k+2)}P_k^{(\alpha,\beta)}(x)P_{n-k}^{(-n-\alpha-1,-n-\beta-1)}(y)\\
&= \sum_{i=0}^{n}(-1)^{n-i}\frac{(\alpha+\beta+1)_{2i}}{i!\,(n-i)!}\left(\frac{x-1}{2}\right)^i\left(\frac{y-1}{2}\right)^{n-i}\frac{\alpha+\beta+2i+1}{\Gamma(\alpha+\beta+2i+2)}\\
&= \frac{1}{\Gamma(\alpha+\beta+1)}\frac{1}{n!}\sum_{i=0}^{n}\binom{n}{i}\left(\frac{x-1}{2}\right)^i\left(\frac{1-y}{2}\right)^{n-i}\\
&= \frac{1}{\Gamma(\alpha+\beta+1)}\frac{1}{n!}\left(\frac{x-y}{2}\right)^n,\ n=0,1,2,\ldots,
\end{aligned}$$

which proves (35).

7 The symmetric generalized ultraspherical polynomials

Let $\alpha>-1$. In [10] we found the coefficients $\{a_i(x)\}_{i=0}^{\infty}$ of the differential equation (32) for the symmetric generalized ultraspherical polynomials $\left\{P_n^{\alpha,\alpha,M,M}(x)\right\}_{n=0}^{\infty}$. In order to do this we had to solve the following two systems of equations for the coefficients $\{a_i(x)\}_{i=0}^{\infty}$:

$$\sum_{i=0}^{\infty} a_i(x)D^i P_n^{(\alpha,\alpha)}(x) = \frac{4}{2\alpha+1}\binom{n+2\alpha}{n} D^2 P_n^{(\alpha,\alpha)}(x) \tag{38}$$

and

$$\sum_{i=0}^{\infty} i a_i(x)D^i P_n^{(\alpha,\alpha)}(x) + x\sum_{i=0}^{\infty} a_i(x)D^{i+1} P_n^{(\alpha,\alpha)}(x) = 4\binom{n+2\alpha+1}{n-1} D^2 P_n^{(\alpha,\alpha)}(x) \tag{39}$$

for $n = 0,1,2,\ldots$, where the coefficients $\{a_i(x)\}_{i=0}^{\infty}$ are continuous functions on the real line and $\{a_i(x)\}_{i=1}^{\infty}$ are independent of n. Now we suppose that $a_0(x) := a_0(n,\alpha)$ is independent of x as we did in [7]. Then it is clear (see for instance lemma 1 in [7]) that $a_i(x)$ must be a polynomial in x of degree at most i for each $i = 1,2,3,\ldots$. In [10] we showed that the solution for $\{a_i(x)\}_{i=0}^{\infty}$ is not unique. In fact it was shown that

$$a_0(x) := a_0(n,\alpha) = a_0(1,\alpha)b_0(n,\alpha) + c_0(n,\alpha),\ n = 0,1,2,\ldots \tag{40}$$

and that

$$a_i(x) = a_0(1,\alpha)b_i(x) + c_i(x),\ i = 1,2,3,\ldots, \tag{41}$$

where $a_0(1,\alpha)$ is arbitrary and

$$b_0(n,\alpha) = \frac{1}{2}\left[1-(-1)^n\right],\ n = 0,1,2,\ldots, \tag{42}$$

$$c_0(n,\alpha) = 4(2\alpha+3)\binom{n+2\alpha+2}{n-2},\ n = 0,1,2,\ldots, \tag{43}$$

$$b_i(x) = \frac{2^{i-1}}{i!}(-x)^i,\ i = 1,2,3,\ldots, \tag{44}$$

$$c_1(x) = 0 \text{ and } c_i(x) = (2\alpha+3)(1-x^2)\frac{2^i}{i!}P_{i-2}^{(\alpha-i+3,\alpha-i+3)}(x),\ i = 2,3,4,\ldots. \tag{45}$$

In this section we will give an alternative proof of this by using the Jacobi inversion formula (31). Note that the special case $\beta = \alpha$ of the Jacobi inversion formula (31) reads

$$\sum_{k=j}^{i} \frac{2\alpha+2k+1}{(2\alpha+k+j+1)_{i-j+1}} \times \\ \times P_{i-k}^{(-\alpha-i-1,-\alpha-i-1)}(x)P_{k-j}^{(\alpha+j,\alpha+j)}(x) = \delta_{ij},\ j \leq i,\ i,j = 0,1,2,\ldots. \tag{46}$$

If we apply this inversion formula to the system of equations

$$\sum_{i=1}^{\infty} A_i(x)D^i P_n^{(\alpha,\alpha)}(x) = F_n(x),\ n = 1,2,3,\ldots, \tag{47}$$

then we find

$$A_i(x) = 2^i \sum_{j=1}^{i} \frac{2\alpha+2j+1}{(2\alpha+j+1)_{i+1}} P_{i-j}^{(-\alpha-i-1,-\alpha-i-1)}(x)F_j(x),\ i = 1,2,3,\ldots. \tag{48}$$

The special case $\beta=\alpha$ of (37) reads

$$\sum_{k=j}^{i}\frac{2\alpha+2k+1}{(2\alpha+k+j+1)_{i-j+1}}\times$$
$$\times P_{i-k}^{(-\alpha-i-1,-\alpha-i-1)}(-x)P_{k-j}^{(\alpha+j,\alpha+j)}(x)=\frac{x^{i-j}}{(i-j)!},\ j\leq i,\ i,j=0,1,2,\ldots. \tag{49}$$

By considering (38) and (39) for $n=0$ and $n=1$ we conclude that $a_0(0,\alpha)=0$, $a_0(1,\alpha)$ is arbitrary and $a_1(x)=-a_0(1,\alpha)x$. For $n=2,3,4,\ldots$ it turns out to be more convenient to use another system of equations instead of (39). By using (20) we find for $i=0,1,2,\ldots$

$$\begin{aligned} & iD^iP_n^{(\alpha,\alpha)}(x)+xD^{i+1}P_n^{(\alpha,\alpha)}(x)=D^i\left[xDP_n^{(\alpha,\alpha)}(x)\right] \\ = \; & nD^iP_n^{(\alpha,\alpha)}(x)+\frac{1}{2}(n+\alpha)D^iP_{n-2}^{(\alpha+1,\alpha+1)}(x),\ n=2,3,4,\ldots. \end{aligned}$$

Combining (38) and (39) we now obtain

$$\begin{aligned} \sum_{i=0}^{\infty}a_i(x)D^iP_{n-2}^{(\alpha+1,\alpha+1)}(x) &= \frac{2}{n+\alpha}\left[4\binom{n+2\alpha+1}{n-1}-\frac{4n}{2\alpha+1}\binom{n+2\alpha}{n}\right]D^2P_n^{(\alpha,\alpha)}(x) \\ &= \frac{8}{n+\alpha}\left[\binom{n+2\alpha+1}{n-1}-\binom{n+2\alpha}{n-1}\right]D^2P_n^{(\alpha,\alpha)}(x) \\ &= \frac{8}{n+\alpha}\binom{n+2\alpha}{n-2}D^2P_n^{(\alpha,\alpha)}(x),\ n=2,3,4,\ldots. \end{aligned}$$

So we conclude that (39) for $n=2,3,4,\ldots$ may be replaced by

$$\sum_{i=0}^{\infty}a_i(x)D^iP_{n-2}^{(\alpha+1,\alpha+1)}(x)=\frac{8}{n+\alpha}\binom{n+2\alpha}{n-2}D^2P_n^{(\alpha,\alpha)}(x),\ n=2,3,4,\ldots. \tag{50}$$

Note that for $n=2$ this implies that $a_0(2,\alpha)=4(2\alpha+3)$.

Since $a_i(x)$ must be a polynomial in x of degree at most i for each $i=1,2,3,\ldots$ we may write

$$a_i(x)=k_ix^i+\text{ lower order terms },\ i=1,2,3,\ldots.$$

By comparing the coefficients of highest degree in (38) and (50) we find by using (8) :

$$\frac{a_0(n,\alpha)}{n!}+\sum_{i=1}^{n}\frac{k_i}{(n-i)!}=0,\ n=1,2,3,\ldots$$

and

$$\frac{a_0(n,\alpha)}{(n-2)!}+\sum_{i=1}^{n-2}\frac{k_i}{(n-i-2)!}=4(2n+2\alpha-1)\binom{n+2\alpha}{n-2}\frac{1}{(n-2)!},\ n=3,4,5,\ldots.$$

Since k_i is independent of n for $i=1,2,3,\ldots$ and $a_0(2,\alpha)=4(2\alpha+3)$ we conclude that

$$a_0(n+2,\alpha)-a_0(n,\alpha)=4(2n+2\alpha+3)\binom{n+2\alpha+2}{n},\ n=0,1,2,\ldots, \tag{51}$$

where $a_0(0,\alpha) = 0$ and $a_0(1,\alpha)$ is arbitrary. Hence

$$a_0(2n,\alpha) - a_0(0,\alpha) = 4\sum_{k=0}^{n-1}\binom{2k+2\alpha+2}{2k}(4k+2\alpha+3),\ n = 1,2,3,\ldots$$

and

$$a_0(2n+1,\alpha) - a_0(1,\alpha) = 4\sum_{k=0}^{n-1}\binom{2k+2\alpha+3}{2k+1}(4k+2\alpha+5),\ n = 1,2,3,\ldots.$$

Since we have by using (23)

$$\begin{aligned}\sum_{k=0}^{n-1}\binom{2k+2\alpha+2}{2k}(4k+2\alpha+3) &= \sum_{k=0}^{n-1}\frac{(2\alpha+3)_{2k+1}}{(2k)!} + \sum_{k=0}^{n-1}\frac{2k(2\alpha+3)_{2k}}{(2k)!}\\ = \sum_{k=0}^{2n-2}\frac{(2\alpha+3)_{k+1}}{k!} &= (2\alpha+3)\frac{(2\alpha+5)_{2n-2}}{(2n-2)!} = (2\alpha+3)\binom{2n+2\alpha+2}{2n-2},\ n = 1,2,3,\ldots\end{aligned}$$

and

$$\begin{aligned}\sum_{k=0}^{n-1}\binom{2k+2\alpha+3}{2k+1}(4k+2\alpha+5) &= \sum_{k=0}^{n-1}\frac{(2\alpha+3)_{2k+2}}{(2k+1)!} + \sum_{k=0}^{n-1}\frac{(2\alpha+3)_{2k+1}}{(2k)!}\\ = \sum_{k=0}^{2n-1}\frac{(2\alpha+3)_{k+1}}{k!} &= (2\alpha+3)\frac{(2\alpha+5)_{2n-1}}{(2n-1)!} = (2\alpha+3)\binom{2n+2\alpha+3}{2n-1},\ n = 1,2,3,\ldots,\end{aligned}$$

we now conclude that (40), (42) and (43) hold.

The systems of equations (38) and (50) lead to

$$\sum_{i=1}^{\infty} a_i(x)D^iP_n^{(\alpha,\alpha)}(x) = \frac{4}{2\alpha+1}\binom{n+2\alpha}{n}D^2P_n^{(\alpha,\alpha)}(x) - a_0(n,\alpha)P_n^{(\alpha,\alpha)}(x) \tag{52}$$

for $n = 0,1,2,\ldots$ and

$$\sum_{i=1}^{\infty} a_i(x)D^iP_{n-2}^{(\alpha+1,\alpha+1)}(x) = \frac{8}{n+\alpha}\binom{n+2\alpha}{n-2}D^2P_n^{(\alpha,\alpha)}(x) - a_0(n,\alpha)P_{n-2}^{(\alpha+1,\alpha+1)}(x) \tag{53}$$

for $n = 2,3,4,\ldots$.

First we remark that (52) is true for $n = 0$ and $n = 1$ since $a_0(0,\alpha) = 0$ and $a_1(x) = -a_0(1,\alpha)x$. Then we will show that every solution of (53) also satisfies (52). Suppose that $\{a_i(x)\}_{i=1}^{\infty}$ is a solution of (53). Now we use (11), (21), (51) and the fact that $\{a_i(x)\}_{i=1}^{\infty}$ are independent of n to obtain for $n = 2,3,4,\ldots$

$$\begin{aligned}&2(n+\alpha+1)(2n+2\alpha+1)\sum_{i=1}^{\infty} a_i(x)D^iP_n^{(\alpha,\alpha)}(x)\\ = \ &(n+2\alpha+1)(n+2\alpha+2)\sum_{i=1}^{\infty} a_i(x)D^iP_n^{(\alpha+1,\alpha+1)}(x) +\\ &\qquad - (n+\alpha)(n+\alpha+1)\sum_{i=1}^{\infty} a_i(x)D^iP_{n-2}^{(\alpha+1,\alpha+1)}(x)\end{aligned}$$

$$\begin{aligned}
&= (n+2\alpha+1)(n+2\alpha+2)\times \\
&\qquad\times\left[\frac{8}{n+\alpha+2}\binom{n+2\alpha+2}{n}D^2P_{n+2}^{(\alpha,\alpha)}(x)-a_0(n+2,\alpha)P_n^{(\alpha+1,\alpha+1)}(x)\right]+ \\
&\qquad-(n+\alpha)(n+\alpha+1)\left[\frac{8}{n+\alpha}\binom{n+2\alpha}{n-2}D^2P_n^{(\alpha,\alpha)}(x)-a_0(n,\alpha)P_{n-2}^{(\alpha+1,\alpha+1)}(x)\right] \\
&= \frac{2(n+2\alpha+1)(n+2\alpha+2)}{n+\alpha+2}\binom{n+2\alpha+2}{n}\times \\
&\qquad\times\Big[(n+2\alpha+3)(n+2\alpha+4)P_n^{(\alpha+2,\alpha+2)}(x)+ \\
&\qquad\qquad -2(n+\alpha+2)(2n+2\alpha+3)P_n^{(\alpha+1,\alpha+1)}(x)\Big]+ \\
&\qquad-8(n+\alpha+1)\binom{n+2\alpha}{n-2}D^2P_n^{(\alpha,\alpha)}(x)+ \\
&\qquad-a_0(n,\alpha)\Big[(n+2\alpha+1)(n+2\alpha+2)P_n^{(\alpha+1,\alpha+1)}(x)+ \\
&\qquad\qquad-(n+\alpha)(n+\alpha+1)P_{n-2}^{(\alpha+1,\alpha+1)}(x)\Big] \\
&= 2(n+\alpha+1)(n+2\alpha+1)(n+2\alpha+2)\binom{n+2\alpha+2}{n}P_{n-2}^{(\alpha+2,\alpha+2)}(x)+ \\
&\qquad-8(n+\alpha+1)\binom{n+2\alpha}{n-2}D^2P_n^{(\alpha,\alpha)}(x)+ \\
&\qquad-2(n+\alpha+1)(2n+2\alpha+1)a_0(n,\alpha)P_n^{(\alpha,\alpha)}(x) \\
&= 8(n+\alpha+1)\left[\binom{n+2\alpha+2}{n}-\binom{n+2\alpha}{n-2}\right]D^2P_n^{(\alpha,\alpha)}(x)+ \\
&\qquad-2(n+\alpha+1)(2n+2\alpha+1)a_0(n,\alpha)P_n^{(\alpha,\alpha)}(x) \\
&= 2(n+\alpha+1)(2n+2\alpha+1)\left[\frac{4}{2\alpha+1}\binom{n+2\alpha}{n}D^2P_n^{(\alpha,\alpha)}(x)-a_0(n,\alpha)P_n^{(\alpha,\alpha)}(x)\right].
\end{aligned}$$

Since $\alpha>-1$ this proves that every solution of (53) also satisfies (52).

Now we will solve (53). Shifting n by two we may write, since the coefficients $\{a_i(x)\}_{i=1}^{\infty}$ are independent of n

$$\sum_{i=1}^{\infty}a_i(x)D^iP_n^{(\alpha+1,\alpha+1)}(x)=F_n(x),\ n=0,1,2,\ldots, \tag{54}$$

where

$$F_n(x)=\frac{8}{n+\alpha+2}\binom{n+2\alpha+2}{n}D^2P_{n+2}^{(\alpha,\alpha)}(x)-a_0(n+2,\alpha)P_n^{(\alpha+1,\alpha+1)}(x).$$

Since $a_0(2,\alpha)=4(2\alpha+3)$ we easily find that $F_0(x)=0$. This implies that the system of equations (54) is of the form (47). So if we apply the inversion formula (46) to the system of equations (54) we obtain by using (48)

$$a_i(x)=2^i\sum_{j=1}^{i}\frac{2\alpha+2j+3}{(2\alpha+j+3)_{i+1}}P_{i-j}^{(-\alpha-i-2,-\alpha-i-2)}(x)F_j(x),\ i=1,2,3,\ldots.$$

Hence, by using (40) we now conclude that the coefficients $\{a_i(x)\}_{i=1}^{\infty}$ can be written in the form (41). Moreover, we find by using (42), (12) and (49)

$$\begin{aligned}
b_i(x) &= 2^{i-1}\sum_{j=1}^{i}\frac{2\alpha+2j+3}{(2\alpha+j+3)_{i+1}}P_{i-j}^{(-\alpha-i-2,-\alpha-i-2)}(x)P_j^{(\alpha+1,\alpha+1)}(x)\left[(-1)^j-1\right]\\
&= 2^{i-1}\sum_{j=0}^{i}\frac{2\alpha+2j+3}{(2\alpha+j+3)_{i+1}}P_{i-j}^{(-\alpha-i-2,-\alpha-i-2)}(x)P_j^{(\alpha+1,\alpha+1)}(-x)+\\
&\qquad -2^{i-1}\sum_{j=0}^{i}\frac{2\alpha+2j+3}{(2\alpha+j+3)_{i+1}}P_{i-j}^{(-\alpha-i-2,-\alpha-i-2)}(x)P_j^{(\alpha+1,\alpha+1)}(x)\\
&= 2^{i-1}\frac{(-x)^i}{i!},\ i=1,2,3,\ldots,
\end{aligned}$$

which proves (44). And by using (43) and (11) we obtain

$$c_i(x)=2^i\sum_{j=1}^{i}\frac{2\alpha+2j+3}{(2\alpha+j+3)_{i+1}}P_{i-j}^{(-\alpha-i-2,-\alpha-i-2)}(x)G_j(x),\ i=1,2,3,\ldots,$$

where

$$G_j(x)=\frac{4(2\alpha+3)}{j+\alpha+2}\binom{j+2\alpha+4}{j}\left[(\alpha+2)P_j^{(\alpha+2,\alpha+2)}(x)-(j+\alpha+2)P_j^{(\alpha+1,\alpha+1)}(x)\right].$$

It is clear that $G_1(x)=0$, which implies that $c_1(x)=0$. Note that since $b_1(x)=-x$ this also implies that $a_1(x)=-a_0(1,\alpha)x$, which agrees with what we have found before. Now we use (22) to find

$$G_j(x)=(2\alpha+3)(1-x^2)\binom{j+2\alpha+4}{j}P_{j-2}^{(\alpha+3,\alpha+3)}(x),\ j=2,3,4,\ldots.$$

Hence, for $i=2,3,4,\ldots$ we have

$$\begin{aligned}
c_i(x) &= (2\alpha+3)(1-x^2)2^i\times\\
&\qquad\times\sum_{j=2}^{i}\frac{2\alpha+2j+3}{(2\alpha+j+3)_{i+1}}\binom{j+2\alpha+4}{j}P_{i-j}^{(-\alpha-i-2,-\alpha-i-2)}(x)P_{j-2}^{(\alpha+3,\alpha+3)}(x).
\end{aligned}$$

Now it remains to show that

$$\begin{aligned}
&\sum_{j=2}^{i}\frac{2\alpha+2j+3}{(2\alpha+j+3)_{i+1}}\binom{j+2\alpha+4}{j}P_{i-j}^{(-\alpha-i-2,-\alpha-i-2)}(x)P_{j-2}^{(\alpha+3,\alpha+3)}(x)\\
&\qquad=\frac{1}{i!}P_{i-2}^{(\alpha-i+3,\alpha-i+3)}(x),\ i=2,3,4,\ldots.
\end{aligned}\tag{55}$$

In order to do this we write for $i=2,3,4,\ldots$

$$\begin{aligned}
&\sum_{j=2}^{i}\frac{2\alpha+2j+3}{(2\alpha+j+3)_{i+1}}\binom{j+2\alpha+4}{j}P_{i-j}^{(-\alpha-i-2,-\alpha-i-2)}(x)P_{j-2}^{(\alpha+3,\alpha+3)}(x)\\
&\qquad=\sum_{k=0}^{i-2}\frac{2\alpha+2k+7}{(2\alpha+k+5)_{i+1}}\frac{(2\alpha+5)_{k+2}}{(k+2)!}P_{i-k-2}^{(-\alpha-i-2,-\alpha-i-2)}(x)P_k^{(\alpha+3,\alpha+3)}(x).
\end{aligned}$$

Now we apply definition (8) to $P_k^{(\alpha+3,\alpha+3)}(x)$ and definition (9) to $P_{i-k-2}^{(-\alpha-i-2,-\alpha-i-2)}(x)$ and change the order of summation to obtain for $i = 2, 3, 4, \ldots$

$$\begin{aligned}
&\sum_{k=0}^{i-2} \frac{2\alpha+2k+7}{(2\alpha+k+5)_{i+1}} \frac{(2\alpha+5)_{k+2}}{(k+2)!} P_{i-k-2}^{(-\alpha-i-2,-\alpha-i-2)}(x) P_k^{(\alpha+3,\alpha+3)}(x) \\
= \;& \sum_{k=0}^{i-2} \sum_{m=0}^{i-k-2} \sum_{n=0}^{k} (-1)^{i-k-2} \times \\
&\qquad \times \frac{(2\alpha+2k+7)(2\alpha+5)_{k+n+2}(\alpha+n+4)_{i-m-n-2}}{(2\alpha+k+5)_{i-m+1}(k+2)!\, m!\, (i-k-m-2)!\, n!\, (k-n)!} \left(\frac{x-1}{2}\right)^{m+n} \\
= \;& \sum_{m=0}^{i-2} \sum_{n=0}^{i-m-2} \sum_{k=0}^{i-m-n-2} (-1)^{i-n-k} \times \\
&\qquad \times \frac{(2\alpha+2n+2k+7)(2\alpha+5)_{2n+k+2}(\alpha+n+4)_{i-m-n-2}}{(2\alpha+n+k+5)_{i-m+1}(n+k+2)!\, m!\, (i-m-n-k-2)!\, n!\, k!} \left(\frac{x-1}{2}\right)^{m+n} \\
= \;& \sum_{j=0}^{i-2} \sum_{n=0}^{j} \sum_{k=0}^{i-j-2} (-1)^{i-n-k} \times \\
&\qquad \times \frac{(2\alpha+2n+2k+7)(2\alpha+5)_{n+k}(\alpha+n+4)_{i-j-2}}{(2\alpha+2n+k+7)_{i-j-1}(n+k+2)!\, (j-n)!\, (i-j-k-2)!\, n!\, k!} \left(\frac{x-1}{2}\right)^{j} \\
= \;& \sum_{j=0}^{i-2} \sum_{n=0}^{j} (-1)^{i-n} \frac{(2\alpha+5)_n(\alpha+n+4)_{i-j-2}}{(2\alpha+2n+7)_{i-j-1}(n+2)!\, (j-n)!\, (i-j-2)!\, n!} \left(\frac{x-1}{2}\right)^{j} \times \\
&\qquad \times \sum_{k=0}^{i-j-2} \frac{(-i+j+2)_k(2\alpha+n+5)_k(2\alpha+2n+7)_k}{(2\alpha+2n+i-j+6)_k(n+3)_k k!} (2\alpha+2n+2k+7).
\end{aligned}$$

Hence, by using (29), (24) and definition (9) we obtain for $i = 2, 3, 4, \ldots$

$$\begin{aligned}
&\sum_{k=0}^{i-2} \frac{2\alpha+2k+7}{(2\alpha+k+5)_{i+1}} \frac{(2\alpha+5)_{k+2}}{(k+2)!} P_{i-k-2}^{(-\alpha-i-2,-\alpha-i-2)}(x) P_k^{(\alpha+3,\alpha+3)}(x) \\
= \;& \sum_{j=0}^{i-2} \sum_{n=0}^{j} (-1)^{i-n} \frac{(2\alpha+5)_n(\alpha+n+4)_{i-j-2}}{(2\alpha+2n+7)_{i-j-1}(n+2)!\, (j-n)!\, (i-j-2)!\, n!} \left(\frac{x-1}{2}\right)^{j} \times \\
&\qquad \times \frac{(2\alpha+2n+7)_{i-j-1}(-\alpha-1)_{i-j-2}}{(n+3)_{i-j-2}(\alpha+n+4)_{i-j-2}} \\
= \;& \sum_{j=0}^{i-2} \sum_{n=0}^{j} (-1)^{i-n} \frac{(2\alpha+5)_n(-\alpha-1)_{i-j-2}}{(n+i-j)!\, (j-n)!\, (i-j-2)!\, n!} \left(\frac{x-1}{2}\right)^{j} \\
= \;& (-1)^i \sum_{j=0}^{i-2} \frac{(-\alpha-1)_{i-j-2}}{(i-j)!\, (i-j-2)!\, j!} \left(\frac{x-1}{2}\right)^{j} {}_2F_1\left(\left.\begin{matrix} -j, 2\alpha+5 \\ i-j+1 \end{matrix}\right| 1\right) \\
= \;& \frac{(-1)^i}{i!} \sum_{j=0}^{i-2} \frac{(-\alpha-1)_{i-j-2}(i-j-2\alpha-4)_j}{(i-j-2)!\, j!} \left(\frac{x-1}{2}\right)^{j} \\
= \;& \frac{(-1)^{i-2}}{i!} \sum_{j=0}^{i-2} \frac{(i-j-2\alpha-4)_j}{j!} \frac{(-\alpha-1)_{i-j-2}}{(i-j-2)!} \left(\frac{x-1}{2}\right)^{j} = \frac{1}{i!} P_{i-2}^{(\alpha-i+3,\alpha-i+3)}(x),
\end{aligned}$$

which proves (55).

8 Some remarks

Note that we have from definition (3) for the Laguerre polynomials that

$$L_n^{(-n)}(x) = (-1)^n \frac{x^n}{n!},\ n = 0, 1, 2, \ldots. \tag{56}$$

Hence, the polynomial $L_n^{(-n)}(x)$ reduces to a monomial of degree n for all $n \in \{0, 1, 2, \ldots\}$. Definition (10) for the Jacobi polynomials leads to

$$P_n^{(-n,\beta)}(x) = \binom{n+\beta}{n} \left(\frac{x-1}{2}\right)^n,\ n = 0, 1, 2, \ldots,$$

which is also a monomial. However, this monomial might reduce to the zero polynomial. For instance, $P_n^{(-n,-n)}(x)$ equals the zero polynomial for all $n \in \{1, 2, 3, \ldots\}$.

Further we remark that if we replace x by $1 - \dfrac{2x}{\beta}$ and y by $1 - \dfrac{2y}{\beta}$ in (35), multiply by $\Gamma(\alpha + \beta + 1)\beta^n$ and let β tend to infinity in an appropriate way we obtain by using (16)

$$\sum_{k=0}^{n} L_k^{(\alpha)}(x) L_{n-k}^{(-n-\alpha-1)}(-y) = \frac{(y-x)^n}{n!},\ n = 0, 1, 2, \ldots. \tag{57}$$

Note that (7) is a special case of (57) since

$$L_{n-k}^{(-n-\alpha-1)}(0) = (-1)^{n-k} \frac{(\alpha+k+1)_{n-k}}{(n-k)!} = (-1)^n (-1)^k \binom{n+\alpha}{n-k},\ k = 0, 1, 2, \ldots, n$$

for all $n \in \{0, 1, 2, \ldots\}$. Moreover, note that (57) is a special case of the well-known convolution formula for the classical Laguerre polynomials

$$\sum_{k=0}^{n} L_k^{(\alpha)}(x) L_{n-k}^{(\beta)}(y) = L_n^{(\alpha+\beta+1)}(x+y),\ n = 0, 1, 2, \ldots$$

in view of (56). By using the technique demonstrated at the end of section 5 this convolution formula can be proved for all α and β which might even depend on n.

We remark that, by using the fact that

$$(b/2)_k (b + 2k) = b(b/2 + 1)_k,\ k = 0, 1, 2, \ldots,$$

the formulas (25) and (29) can also be obtained by using summation formulas for terminating well-poised hypergeometric series (see for instance formulas (III.9) and (III.26) in [15]).

The computation of $a_0(n, \alpha)$ from (51) can also be done in the following way. Note that we have

$$(2n + 2\alpha + 3)\binom{n+2\alpha+2}{n} = (2\alpha + 3)\left[\binom{n+2\alpha+4}{n} - \binom{n+2\alpha+2}{n-2}\right],\ n = 0, 1, 2, \ldots.$$

Hence, instead of (23) we can also use the telescoping property of the sums to find that

$$\sum_{k=0}^{n-1} \binom{2k+2\alpha+2}{2k}(4k+2\alpha+3) = (2\alpha+3)\binom{2n+2\alpha+2}{2n-2},\ n = 1,2,3,\ldots$$

and

$$\sum_{k=0}^{n-1} \binom{2k+2\alpha+3}{2k+1}(4k+2\alpha+5) = (2\alpha+3)\binom{2n+2\alpha+3}{2n-1},\ n = 1,2,3,\ldots.$$

Finally, we remark that if we apply the inversion formula (46) to the system of equations (52) instead of (53) we find for $i = 1,2,3,\ldots$ that

$$b_i(x) = 2^{i-1}\sum_{j=1}^{i} \frac{2\alpha+2j+1}{(2\alpha+j+1)_{i+1}} P_{i-j}^{(-\alpha-i-1,-\alpha-i-1)}(x) P_j^{(\alpha,\alpha)}(x)\left[(-1)^j - 1\right] \tag{58}$$

and

$$\begin{aligned} c_i(x) &= 2^{i+2}\sum_{j=1}^{i} \frac{2\alpha+2j+1}{(2\alpha+j+1)_{i+1}} P_{i-j}^{(-\alpha-i-1,-\alpha-i-1)}(x) \times \\ &\times \left[\frac{1}{2\alpha+1}\binom{j+2\alpha}{j} D^2 P_j^{(\alpha,\alpha)}(x) - (2\alpha+3)\binom{j+2\alpha+2}{j-2} P_j^{(\alpha,\alpha)}(x)\right]. \end{aligned} \tag{59}$$

From (58) we easily obtain (44) in the same way as before by using (12), (46) and (49). Further we easily find from (59) that $c_1(x) = 0$, but we were not able to derive (45) for $i = 2,3,4,\ldots$ from (59).

References

[1] H. Bavinck : *A direct approach to Koekoek's differential equation for generalized Laguerre polynomials.* Acta Mathematica Hungarica **66**, 1995, 247-253.

[2] H. Bavinck : *A difference operator of infinite order with Sobolev-type Charlier polynomials as eigenfunctions.* Indagationes Mathematicae **7**, 1996, 281-291.

[3] H. Bavinck & H. van Haeringen : *Difference equations for generalized Meixner polynomials.* Journal of Mathematical Analysis and Applications **184**, 1994, 453-463.

[4] H. Bavinck & R. Koekoek : *On a difference equation for generalizations of Charlier polynomials.* Journal of Approximation Theory **81**, 1995, 195-206.

[5] T.S. Chihara : *An introduction to orthogonal polynomials.* Mathematics and Its Applications **13**, Gordon and Breach, New York, 1978.

[6] J. Koekoek & R. Koekoek : *On a differential equation for Koornwinder's generalized Laguerre polynomials.* Proceedings of the American Mathematical Society **112**, 1991, 1045-1054.

[7] J. Koekoek, R. Koekoek & H. Bavinck : *On differential equations for Sobolev-type Laguerre polynomials.* Transactions of the American Mathematical Society, to appear.

[8] R. Koekoek : *Generalizations of the classical Laguerre polynomials and some q-analogues.* Delft University of Technology, Thesis, 1990.

[9] R. Koekoek : *The search for differential equations for certain sets of orthogonal polynomials.* Journal of Computational and Applied Mathematics **49**, 1993, 111-119.

[10] R. Koekoek : *Differential equations for symmetric generalized ultraspherical polynomials.* Transactions of the American Mathematical Society **345**, 1994, 47-72.

[11] R. Koekoek & H.G. Meijer : *A generalization of Laguerre polynomials.* SIAM Journal on Mathematical Analysis **24**, 1993, 768-782.

[12] R. Koekoek & R.F. Swarttouw : *The Askey-scheme of hypergeometric orthogonal polynomials and its q-analogue.* Delft University of Technology, Faculty of Technical Mathematics and Informatics, report no. **94-05**, 1994.

[13] T.H. Koornwinder : *Orthogonal polynomials with weight function* $(1-x)^{\alpha}(1+x)^{\beta}+M\delta(x+1)+N\delta(x-1)$. Canadian Mathematical Bulletin **27**(2), 1984, 205-214.

[14] Y.L. Luke : *The special functions and their approximations.* Volume I. Academic Press, San Diego, 1969.

[15] L.J. Slater : *Generalized hypergeometric functions.* Cambridge University Press, Cambridge, 1966.

[16] G. Szegö : *Orthogonal polynomials.* American Mathematical Society Colloquium Publications **23** (1939), Fourth edition, Providence, Rhode Island, 1975.

Menelaoslaan 4, 5631 LN Eindhoven, The Netherlands

Delft University of Technology, Faculty of Technical Mathematics and Informatics,
P.O. Box 5031, 2600 GA Delft, The Netherlands, e-mail : koekoek@twi.tudelft.nl